

Table 8-2. Hypersensitivity pneumonitis: Mortality rates (per million population) by race and sex, U.S. residents age 15 and over, 1990-1999

Year	Overall	White		Black		Other	
		Male	Female	Male	Female	Male	Female
Crude Mortality Rate							
1990	0.21	0.36	0.11	0.10	—	0.29	0.27
1991	0.18	0.32	0.09	—	—	0.55	—
1992	0.09	0.16	0.05	0.09	—	—	—
1993	0.23	0.38	0.15	—	0.08	—	0.24
1994	0.18	0.34	0.09	—	—	—	—
1995	0.18	0.26	0.11	0.18	0.23	—	—
1996	0.25	0.40	0.17	0.09	0.08	—	—
1997	0.18	0.29	0.14	—	—	—	—
1998	0.18	0.35	0.08	0.08	—	—	—
1999	0.27	0.40	0.23	0.08	—	—	—
1990-1999	0.19	0.33	0.12	0.06	0.04	0.07	0.04
Age-Adjusted Mortality Rate							
1990	0.22	0.41	0.09	0.07	—	0.42	0.34
1991	0.19	0.40	0.08	—	—	0.95	—
1992	0.10	0.18	0.05	0.20	—	—	—
1993	0.24	0.45	0.13	—	0.11	—	0.54
1994	0.19	0.43	0.08	—	—	—	—
1995	0.19	0.33	0.10	0.31	0.28	—	—
1996	0.25	0.46	0.14	0.07	0.09	—	—
1997	0.19	0.35	0.12	—	—	—	—
1998	0.18	0.40	0.06	0.10	—	—	—
1999	0.27	0.46	0.20	0.10	—	—	—
1990-1999	0.20	0.39	0.11	0.09	0.05	0.12	0.08

— indicates no deaths listed.

NOTE: See appendices for source description, methods, and ICD codes.

SOURCE: National Center for Health Statistics multiple cause of death data. Population estimates from U.S. Bureau of the Census.